
Testicular Cancer Reference Handout



Testicular Cancer Tx Stage I-IIA

SEMINOMA NO elevation in tumor markers NO Poor Risk Group

Orchiectomy +			
Stage I NO	Observation Consider RT OR Consider chemo		
Stage II N+	Adjuvant RT OR Adjuvant Chemo (Carbo x 1-2C)		
	Stage I NO Stage II		

NO RPLND for seminomas!

NON-SEMINOMAYES elevation in tumor markersMIXEDYES Poor Risk Group

Orchiectomy +				
Stage I NO	Observation Consider RLPND OR Consider chemo (BEP x 1-2C)			
Stage II N+	Adjuvant RPLND OR Adjuvant Chemo (BEP x1-2C)			

NO RT for non-seminomas!

	Testicular Cancer Tx Stage IIB-IV									
\square	SEMINOMA					NON-SEMINOMA/MIXED SEMINOMA				
NO elevation in tumor markers NO Poor Risk Group				YES elevation in tumor markers YES Poor Risk Group						
	Good Risk		BEP x3 EP x4			Good Risk	BEP x3 EP x4			
	Intermediate Risk		BEP x4 VIP x4			Intermediate Risk Poor Risk	BEP x4 VIP x4			
Chemo Regimens										
	BEP: (B) Bleomycin (E) Etoposide (P) Cisplatin			C	Chemo Side Effects					
	EP: (E) Etoposide (P) Cisplatin		Ci Ifo	Bleomycin → Pneumonitis, Rash Cisplatin → Neuropathy, nephropathy, ototoxicity Ifosfamide → Hemorrhagic cystitis, CNS toxicity Paclitaxel → Neuropathy						
	VIP:	(V) Etoposid (I) Ifosfamic (P) Cisplatir	le							